

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 July 2001 (26.07.2001)

PCT

(10) International Publication Number
WO 01/053452 A3

(51) International Patent Classification⁷: C12N 15/82, A01H 5/00, C12N 5/10

(21) International Application Number: PCT/SG00/00010

(22) International Filing Date: 19 January 2000 (19.01.2000)

(25) Filing Language: English

(26) Publication Language: English

(71) Applicant (for all designated States except US): **INSTITUTE OF MOLECULAR AGROBIOLOGY [SG/SG]**; 1 research Link, The National University of Singapore, Singapore 117604 (SG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **XIE, Deyu [CN/SG]**; Blk 702, West Coast Road, #09-535, Singapore 120702 (SG). **HONG, Yan [SG/SG]**; Blk 206, 21 Bukit Batok Street, #06-80, Singapore 650206 (SG).

(74) Agent: **MIRANDAH, Gladys**; Ella Cheong & G. Mirandah, Raffles City, P.O. Box 0931, Singapore 911732 (SG).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:


— with international search report

(88) Date of publication of the international search report:
7 November 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: REGENERATION AND GENETIC TRANSFORMATION OF *ACACIA MANGIUM*

Adventitious bud induction from leaf
A: Callus induction; B: Adventitious bud recovery;
C,D,E: Plumule formation and adventitious bud elongation



(57) Abstract: The present invention is directed to a method of *Acacia mangium* regeneration through organogenesis and a method of genetic transformation of *Acacia mangium*. The method of regeneration comprises inducing callus from different parts of seedlings and vegetatively micropropagated plantlets as explants; adventitious bud induction followed by pinnate leaf and bud elongation and eventually, elongated shoots were induced to root. Based on the regeneration system, a marker gene GUS under cauliflower mosaic virus promoter was introduced to *Acacia mangium* via *Agrobacterium* infection. GUS staining in the regenerated plants and Southern blot hybridization prove the incorporation of the foreign gene into the host genome and expression of the foreign gene.

TECH CENTER 1600/2900

JAN 03 2003

RECEIVED

WO 01/053452 A3

INTERNATIONAL SEARCH REPORT

Inte val Application No

PC 1/56 00/00010

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/82 A01H5/00 C12N5/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01H C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS, EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section PQ, Week 199551 Derwent Publications Ltd., London, GB; Class P13, AN 1995-395221 XP002185994 & JP 07 255304 A (OJI PAPER CO), 9 October 1995 (1995-10-09) abstract	1-6
X	----- GALIANA A ET AL: "IN-VITRO PROPAGATION OF THE NITROGEN-FIXING TREE-LEGUME ACACIA-MANGIUM WILLD" PLANT AND SOIL, vol. 135, no. 2, 1991, pages 151-160, XP001034851 ISSN: 0032-079X the whole document ----- -/-	11-13

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

18 December 2001

Date of mailing of the international search report

09/01/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Holtorf, S

INTERNATIONAL SEARCH REPORT

International Application No

PCT/SG 00/00010

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>BHASKAR P ET AL: "Micropropagation of Acacia mangium Willd through nodal bud culture." INDIAN JOURNAL OF EXPERIMENTAL BIOLOGY, vol. 34, no. 6, 1996, pages 590-591, XP001050518 ISSN: 0019-5189 the whole document</p> <p>---</p>	11
X	<p>KAUR K ET AL: "Plants obtained from the Khair tree (Acacia catechu Willd.) using mature nodal segments." PLANT CELL REPORTS, vol. 17, no. 5, March 1998 (1998-03), pages 427-429, XP001034850 ISSN: 0721-7714 the whole document</p> <p>---</p>	11-13
X	<p>ROUT G R ET AL: "Somatic embryogenesis and plant regeneration from callus culture of Acacia catechu - a multipurpose leguminous tree." PLANT CELL TISSUE AND ORGAN CULTURE, vol. 42, no. 3, 1995, pages 283-285, XP000989635 ISSN: 0167-6857 the whole document</p> <p>---</p>	1-4
X	<p>FAKHRAI H ET AL: "IN-VITRO CULTURE AND PLANT REGENERATION IN VICIA-FABA-SSP-EQUINA CULTIVAR SPRING BLAZE" JOURNAL OF EXPERIMENTAL BOTANY, vol. 40, no. 216, 1989, pages 813-818, XP001034860 ISSN: 0022-0957 the whole document</p> <p>---</p>	25-27
A	<p>MURTHY B N S ET AL: "Thidiazuron: A potent regulator of in vitro plant morphogenesis." IN VITRO CELLULAR & DEVELOPMENTAL BIOLOGY PLANT, vol. 34, no. 4, October 1998 (1998-10), pages 267-275, XP000989563 ISSN: 1054-5476 abstract; page 269, last sentence; page 270, left column;</p> <p>---</p> <p style="text-align: center;">-/--</p>	

INTERNATIONAL SEARCH REPORT

Inte ial Application No

PC 1, 3G 00/00010

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	HUETTEMAN CARL A ET AL: "Thidiazuron: A potent cytokinin for woody plant tissue culture." PLANT CELL TISSUE AND ORGAN CULTURE, vol. 33, no. 2, 1993, pages 105-119, XP000989639 ISSN: 0167-6857 the whole document ---	
A	WO 97 23126 A (EDWARDS GLYN ALYN ;PURSE JOHN GILBERT (GB); SHELL INT RESEARCH (NL) 3 July 1997 (1997-07-03) example 11 ---	
T	XIE DEYU ET AL: "In vitro regeneration of Acacia mangium via organogenesis." PLANT CELL TISSUE AND ORGAN CULTURE, vol. 66, no. 3, 2001, pages 167-173, XP001034852 ISSN: 0167-6857 the whole document -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/SG 09/00010

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
JP 7255304	A	09-10-1995	NONE	
WO 9723126	A	03-07-1997	AT 204120 T	15-09-2001
			AU 721423 B2	06-07-2000
			AU 1303197 A	17-07-1997
			BR 9612217 A	05-10-1999
			CN 1219102 A	09-06-1999
			DE 69614569 D1	20-09-2001
			WO 9723126 A2	03-07-1997
			EP 0877546 A2	18-11-1998
			HU 9901292 A2	28-07-1999
			JP 2000510325 T	15-08-2000

PATENT COOPERATION TREATY

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference GM/MC/R33-71	FOR FURTHER ACTION <small>see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.</small>	
International application No. PCT/SG 00/00010	International filing date (day/month/year) 19/01/2000	(Earliest) Priority Date (day/month/year)
Applicant INSTITUTE OF MOLECULAR AGROBIOLOGY		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 5 sheets.
☐ It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.
- ☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).
- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :
- ☐ contained in the international application in written form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☐ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,

- ☒ the text is approved as submitted by the applicant.
- ☐ the text has been established by this Authority to read as follows:


5. With regard to the **abstract**,

- ☒ the text is approved as submitted by the applicant.
- ☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.

- ☒ as suggested by the applicant.
- ☐ because the applicant failed to suggest a figure.
- ☐ because this figure better characterizes the invention.

1
☐ None of the figures.



INTERNATIONAL SEARCH REPORT

International Application No
PCT/SG 00/00010

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/82 A01H5/00 C12N5/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A01H C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
BIOSIS, EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section PQ, Week 199551 Derwent Publications Ltd., London, GB; Class P13, AN 1995-395221 XP002185994 & JP 07 255304 A (OJI PAPER CO), 9 October 1995 (1995-10-09) abstract	1-6
X	GALIANA A ET AL: "IN-VITRO PROPAGATION OF THE NITROGEN-FIXING TREE-LEGUME ACACIA-MANGIUM WILLD" PLANT AND SOIL, vol. 135, no. 2, 1991, pages 151-160, XP001034851 ISSN: 0032-079X the whole document	11-13
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

18 December 2001

Date of mailing of the international search report

09/01/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Holtorf, S


INTERNATIONAL SEARCH REPORT

International Application No

PCT/SG 00/00010

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>BHASKAR P ET AL: "Micropropagation of Acacia mangium Willd through nodal bud culture." INDIAN JOURNAL OF EXPERIMENTAL BIOLOGY, vol. 34, no. 6, 1996, pages 590-591, XP001050518 ISSN: 0019-5189 the whole document</p> <p>---</p>	11
X	<p>KAUR K ET AL: "Plants obtained from the Khair tree (Acacia catechu Willd.) using mature nodal segments." PLANT CELL REPORTS, vol. 17, no. 5, March 1998 (1998-03), pages 427-429, XP001034850 ISSN: 0721-7714 the whole document</p> <p>---</p>	11-13
X	<p>ROUT G R ET AL: "Somatic embryogenesis and plant regeneration from callus culture of Acacia catechu - a multipurpose leguminous tree." PLANT CELL TISSUE AND ORGAN CULTURE, vol. 42, no. 3, 1995, pages 283-285, XP000989635 ISSN: 0167-6857 the whole document</p> <p>---</p>	1-4
X	<p>FAKHRAI H ET AL: "IN-VITRO CULTURE AND PLANT REGENERATION IN VICIA-FABA-SSP-EQUINA CULTIVAR SPRING BLAZE" JOURNAL OF EXPERIMENTAL BOTANY, vol. 40, no. 216, 1989, pages 813-818, XP001034860 ISSN: 0022-0957 the whole document</p> <p>---</p>	25-27
A	<p>MURTHY B N S ET AL: "Thidiazuron: A potent regulator of in vitro plant morphogenesis." IN VITRO CELLULAR & DEVELOPMENTAL BIOLOGY PLANT, vol. 34, no. 4, October 1998 (1998-10), pages 267-275, XP000989563 ISSN: 1054-5476 abstract; page 269, last sentence; page 270, left column;</p> <p>---</p> <p style="text-align: center;">-/--</p>	




INTERNATIONAL SEARCH REPORT

International Application No

PCT/SG 00/00010

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>HUETTEMAN CARL A ET AL: "Thidiazuron: A potent cytokinin for woody plant tissue culture."</p> <p>PLANT CELL TISSUE AND ORGAN CULTURE, vol. 33, no. 2, 1993, pages 105-119, XP000989639</p> <p>ISSN: 0167-6857</p> <p>the whole document</p>	
A	<p>WO 97 23126 A (EDWARDS GLYN ALYN ;PURSE JOHN GILBERT (GB); SHELL INT RESEARCH (NL) 3 July 1997 (1997-07-03) example 11</p>	
T	<p>XIE DEYU ET AL: "In vitro regeneration of Acacia mangium via organogenesis."</p> <p>PLANT CELL TISSUE AND ORGAN CULTURE, vol. 66, no. 3, 2001, pages 167-173, XP001034852</p> <p>ISSN: 0167-6857</p> <p>the whole document</p>	




INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No


PCT/SG 00/00010

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
JP 7255304	A	09-10-1995	NONE	
WO 9723126	A	03-07-1997	AT 204120 T	15-09-2001
			AU 721423 B2	06-07-2000
			AU 1303197 A	17-07-1997
			BR 9612217 A	05-10-1999
			CN 1219102 A	09-06-1999
			DE 69614569 D1	20-09-2001
			WO 9723126 A2	03-07-1997
			EP 0877546 A2	18-11-1998
			HU 9901292 A2	28-07-1999
			JP 2000510325 T	15-08-2000



XP-002185994

AN - 1995-395221 [04]
AP - JP19940071294 19940317
CPY - OJIP
DC - P13
FS - GMPI
IC - A01H4/00
PA - (OJIP) OJI PAPER CO
PN - JP7255304 A 19951009 DW199551 A01H4/00 009pp
PR - JP19940071294 19940317
XIC - A01H-004/00
AB - J07255304 The proliferation method is applicable to medium which is a combination of cyatokainine system type phytohormone in 0.01mg - 10mg and auxin system type phytohormone in 0 - 10mg.
- ADVANTAGE - Applies to gene technique and organization culture of acacia species.
- (Dwg.0/0)
IW - ACACIA PLANT PROLIFERATION METHOD AUXIN SYSTEM SPECIFIED AMOUNT RANGING
IKW - ACACIA PLANT PROLIFERATION METHOD AUXIN SYSTEM SPECIFIED AMOUNT RANGING
NC - 001
OPD - 1994-03-17
ORD - 1995-10-09
PAW - (OJIP) OJI PAPER CO
TI - Acacia plant proliferation method - by using phytohormone of auxin system by specified amount ranging between 0.01mg and 10mg



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 July 2001 (26.07.2001)


PCT

(10) International Publication Number
WO 01/53452 A2

- (51) International Patent Classification⁷: C12N
- (21) International Application Number: PCT/SG00/00010
- (22) International Filing Date: 19 January 2000 (19.01.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (71) Applicant (for all designated States except US): **INSTITUTE OF MOLECULAR AGROBIOLOGY [SG/SG]**; 1 research Link, The National University of Singapore, Singapore 117604 (SG).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **XIE, Deyu [CN/SG]**; Blk 702, West Coast Road, #09-535, Singapore 120702 (SG); **HONG, Yan [SG/SG]**; Blk 206, 21 Bukit Batok Street, #06-80, Singapore 650206 (SG).
- (74) Agent: **MIRANDAH, Gladys; Ella Cheong & G. Mirandah, Raffles City, P.O. Box 0931, Singapore 911732 (SG).**
- (81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.


(54) Title: REGENERATION AND GENETIC TRANSFORMATION OF *ACACIA MANGIUM*

Adventitious bud induction from leaf
A: Callus induction; B: Adventitious bud recovery;
C,D,E: Plumule formation and adventitious bud elongation



(57) Abstract: The present invention is directed to a method of *Acacia mangium* regeneration through organogenesis and a method of genetic transformation of *Acacia mangium*. The method of regeneration comprises inducing callus from different parts of seedlings and vegetatively micropropagated plantlets as explants; adventitious bud induction followed by pinnate leaf and bud elongation and eventually, elongated shoots were induced to root. Based on the regeneration system, a marker gene GUS under cauliflower mosaic virus promoter was introduced to *Acacia mangium* via *Agrobacterium* infection. GUS staining in the regenerated plants and Southern blot hybridization prove the incorporation of the foreign gene into the host genome and expression of the foreign gene.

WO 01/53452 A2



TITLE OF THE INVENTIONREGENERATION AND GENETIC TRANSFORMATION OF *ACACIA MANGIUM*BACKGROUND OF THE INVENTION

Forests are very important to the world economy and for maintaining and preserving our ecosystem. Forest trees have a wide range of commercial uses (timber for construction, raw material for paper and pulp production, and as an energy source). The global demand for wood products (mostly for paper and pulp and firewood in developing countries) has been increasing year by year when the natural forests are in short supply. Re-forestation is the solution to meeting such increasing demand. Usually, the fast-growing, widely adapted tree species are chosen for re-forestation. Most tree-improvement programs are based on the management of genetic resources, including the selection of superior clones from existing forests, the conservation of genetic variability, partially controlled propagation and classical breeding for desired traits. In spite of the fact that it usually takes several generations to breed, this traditional breeding has been successful in getting elite trees with fast and uniform growth. However, many other traits such as disease and insect resistance, different lignin composition and content are difficult to acquire mainly due to high heterozygosity in tree species and big segregation population. Moreover, the gene(s) conferring certain phenotypes like disease resistance may not be in the gene pool at all. On the other hand, molecular breeding based on genetic transformation of tree species offers the possibility to introduce a particular phenotype without affecting the genetic background of a cultivar. Genetic transformation in *Populus* species and *Eucalyptus* species enabled some success in modification of lignin content (Tzfira et al., 1998; Robinson, 1999). The precondition of molecular breeding of forest tree species is the availability of a reliable and reproducible genetic transformation method, which in turn relies on a system of regeneration of one whole plant from a single cell.

Genus *Acacia* comprises about 1200 tropical and subtropical tree species. It belongs to the family *Mimosaceae*. *Acacia mangium* is a multipurpose, fast growing and nitrogen fixing elite tropical legume tree. An adult tree is up to 30-metres tall and its bole (trunk) is often straight to over half the total height. The true leaf of a seedling is a fern-like pinnate leaf. The first pinnate leaf of a seedling has 6 or 8 leaflets, and then the bipinnate leaf develops from the second pinnate leaf on a seedling. Usually when the young seedling grows up to 8-12 bipinnate leaves, the petiole is dilated into a phyllode, while the leaflets abort completely and the true leaf disappears

from the young seedling. Phyllodes are flattened leaf stems which look and act like the ordinary leaves of other plants. Branchlets, phyllodes and petioles are glabrous or slightly scurfy. Phyllodes are 5-10 cm broad, 2-4 times as long as broad, dark green, and are chartaceous when dry. The phyllodes have 3-4 longitudinal main nerves which join on the dorsal margin at the base of the phyllode. Secondary nerves are fine and inconspicuous.

Flowers are in loose spikes to 10 cm long and are solitary or paired in the upper axils. Flowers are pentamerous and the calyx is 0.6-0.8 mm long with short obtuse lobes. The corolla is twice as long as the calyx. Pods are linear, glabrous, 3-5 mm broad, about 7.5 cm long when green, woody, coiled and brackish-brown when mature, and depressed between the seeds. Seeds are lustrous, black, ellipsoid, ovate or oblong, 3.5×2.5 mm, with the orangish funicle forming a fleshy aril beneath the seed (Duke, 1984). Due to *A. mangium* fast growth, tolerance of infertile soil and high quality fibre, it has been increasingly used for reforestation plantation and soil rehabilitation in degraded soil. Its plantation has been studied for many years in many countries or regions, especially in tropical regions and subtropical regions, such as Australian, Indonesia, Malaysia, India, Thailand, Hawaii, China and Taiwan. Many *A. mangium* plantations have been established in acidic soil or abandoned land or Imperata grassland, for example, in Bangladesh (Latif et al., 1995); in Sabah (Latif et al., 1995; Williams et al., 1992) and in Serdong (Majid et al., 1994; Awang, 1994) in Malaysia; Sangmelina, Cameroon, Kenya (Duguma et al., 1994); Skaerat, Thailand (Khemnark, 1994); Hawaii, USA (Cole et al., 1996); Bogor (Anwar, 1992; Wibowo et al., 1992), Paseh and Kadipaten (Widiarti and Alrasjid, 1987) in Indonesia; Bengal, India (Basu et al., 1987), etc.

Indonesia, with some of the world's largest paper and pulp mills, has been increasingly relying on plantations as the source of wood and *A. mangium* is the preferred choice. Asia Paper and Pulp group has two affiliate companies with a total concession of 540,000 hectares. By 1996, one company had planted 123,000 hectares of *A. mangium*, about 90% of all its plantation, which represented 180 million seedlings. It is estimated that by 2004, Asia Paper and Pulp group will virtually source all its wood from plantation, mainly *A. mangium* plantation (Bayliss, 1998a; Bayliss, 1998b).

Besides its use for paper and pulp, *A. mangium* timber can be used for other applications such as cement banded particleboard, plywood and decorative panel manufacture (Yusoff et al., 1989; Wong et al., 1988).

A large scale of mono-plantation of vegetative propagation from a single plant faces a higher risk of disease infection. It has been found that diseases spread very fast in this kind of plantation and cause large economic losses. Many diseases devastate *A. mangium*: Cinnamon fungus (*Phytophthora cinnamomi*) infection results in collapse and death of plants; Seedling blight, defoliation and dieback due to *Glomerella cingulata* caused serious losses to *A. mangium* in nurseries. Leaf spot by *Cyclindrocladium quinqueseptatum* causes defoliation of seedlings and young trees; Powdery mildew (*Oidium spp*) severely affected *A. mangium* seedlings in nurseries in Thailand; Red rot diseases, caused by *ganoderma sp.* affects *A. mangium* in Malaysia; Brown root disease caused by *Phellinus noxium* affects *A. mangium* in Malaysia and the Solomon islands (Simmons, 1987; Gutteridge and Shelton, 1994). Traditional breeding has been less successful in getting disease resistance in *A. mangium*, mostly likely due to lack of such a gene in the natural gene pool. Molecular breeding by introducing foreign disease resistance genes becomes an important option.

Studies on *A. mangium* tissue culture have been limited to micropropagation (Bhaskan and Subbash, 1996; Ahmad, 1991; Galiana et al., 1991a; Galiana et al., 1991b). A combination of traditional breeding of elite trees and techniques of large-scale vegetative propagation make the large-scale plantation possible. There has been no report on regeneration or genetic transformation of *A. mangium*. Our invention describes the conditions for regeneration of whole plant via organogenesis and a genetic transformation system.

The publications and other materials used herein to illuminate the background of the invention or provide additional details respecting the practice, are incorporated by reference, and for convenience are respectively grouped in the appended List of References.

SUMMARY OF THE INVENTION

A. System of regeneration from callus in *Acacia mangium*

Different parts of *Acacia mangium* seedlings or vegetative micropropagated plantlets (hypocotyl, cotyledon, leaflet, petiole and stem) were used as explants. The seedlings were from isolated seed embryo culture and the plantlets were obtained by micropropagation from meristem regions of a 2-year old tree. Callus formation was induced by MS basic medium (Murashige and Skoog, 1962) supplemented with auxins (either 2,4-D from 0.5 mg/L to 5 mg/L or α -

naphthaleneacetic acid (NAA) from 0.5 mg/L to 2 mg/L) and cytokinins (either kinetin (KT) 0.5 mg/L to 3.0 mg/L or 6-benzylaminopurine (6-BA or BA) 0.5 mg/L to 3.0 mg/L).

Adventitious bud induction was achieved on medium AM-265 (MS basic medium with thidiazuron (1-phenyl-3-(1,2,3-thiadiazol-5-yl)urea or TDZ) 1.0 mg/L, indole-3-acetic acid (IAA) 0.25 mg/L, casein enzymatic hydrolysate (CH) 100 mg/L, L-ascorbic acid (vitamin C or Vc) 100 mg/L, L-glutamine (Gln) 150 mg/L, L-asparagine monohydrate (Asn) 150 mg/L, L-proline (Pro) 150 mg/L, pH 5.8 after autoclaving, phytigel 0.275 or 0.30%, or agar 0.8%, sucrose 30 g/L) under the growth conditions of 16/8 hours light/dark (L/D) cycle and at 28°C.

Pinnate leaf formation and adventitious bud elongation were obtained on medium AM-337 (MS basic medium with TDZ 0.01 mg/L, CH 100 mg/L, Vc 100 mg/L, Gln 150 mg/L, Asn 150 mg/L and Pro 150 mg/L, pH 5.8 after autoclaving, phytigel 0.275 or 0.30%, or agar 0.8%, sucrose 30 g/L), or AM-41 (MS basic medium supplemented with 6-BA 2 mg/L, CH 100 mg/L, Vc 100 mg/L, Gln 150 mg/L, Asn 150 mg/L and Pro 150 mg/L, pH 5.8 after autoclaving, phytigel 0.275 or 0.30%, or agar 0.8%, sucrose 30 g/L) supplemented with gibberellic acid (GA₃) 0.5-2.5 mg/L. The growth conditions are 16/8 hrs (L/D) cycle and at 28°C.

Adventitious buds formed roots and complete plantlets were obtained on MS or ½MS basic medium supplemented with NAA 0.5 to 3.0 mg/L combined with KT 0-0.5 mg/L or IBA (indole butyric acid) 0.01-0.5 mg/L and CH 100 mg/L, Vc 100 mg/L, Gln 150 mg/L, Asn 150 mg/L and Pro 150 mg/L, pH 5.8 after autoclaving, phytigel 0.35%, or agar 1.4%, sucrose 20 or 30 g/L under the growth conditions of 16/8 hrs (L/D) cycle and at 28°C. Complete plantlets were transplanted into peat soil:white sand (3:1) in a growth chamber at a photoperiod of 16/8 hrs (L/D) and 25°C.

B. System of genetic transformation of *A. mangium*

Based on the above established regeneration system, *Agrobacterium* strain LBA4404 (Ooms et al., 1981) harbouring binary vector pBI121 in which a GUS gene is under regulation of an 800 bp cauliflower mosaic virus promoter was used for genetic transformation of *A. mangium*. The *Agrobacteria* with the plasmid were co-cultured with the following explants: (1) Young adventitious shoots vegetatively micropropagated from auxiliary buds dissected from a 2-year old tree. The shoots consisted of young stem and pinnate leaves but no phyllodes. Stem pieces of about 2-3 cm in length after separating pinnate leaf and auxiliary buds were used as

explants. (2) Petiole pieces of 1-2 cm in length and leaflet pieces of 0.4-0.5×0.3-0.4 (cm) from seedlings derived from embryo culture were used as explants. The explants were pre-cultured on AM-265 for three days at 16/8 hrs (L/D) cycle and at 28°C before infection with activated *Agrobacterium* for 15 minutes. Infected explants were then cultured for 3 days in the same media in the dark at 22°C. They were then washed and put into AM-265 to induce callus and adventitious buds. Transgenic plants were selected with G418 in media in four stages: G418 12 mg/L for 25 days, G418 20 mg/L for 60 days, G418 30 mg/L for 25 days and then on G418 12 mg/L. Adventitious bud elongation was induced by AM-265 with GA₃ 2.5 to 5 mg/L in the presence of G418 12 mg/L. For pinnate leaf formation, adventitious buds were transferred into AM-41 with GA₃ 2.5 mg/L at growth conditions of 16/8 hours (L/D) cycle and 28°C. Regenerated shoots were excised and checked for GUS staining and 16% tested positive. Putative transgenic adventitious buds were transferred into AM-357 or AM-451 with or without G418 (10 mg/L) at the photoperiod of 16/8 hours (L/D) to induce rooting. Southern blot hybridization showed positive results when NPTII gene was used as probe.

BRIEF DESCRIPTION OF THE FIGURES

Figures 1A-E show adventitious bud induction from leaf. Figure 1A shows callus induction. Figure 1B shows adventitious bud recovery. Figures 1C, 1D and 1E show plumule formation and adventitious bud elongation.

Figures 2A-C show *Acacia mangium* regeneration from petiole. Figure 2C shows callus induction. Figure 2B shows adventitious bud recovery. Figure 2C shows plumule formation and adventitious bud elongation.

Figures 3A-E demonstrate *Acacia mangium* regeneration from stem pieces. Figure 3A shows regeneration from callus. Figure 3B shows adventitious buds induced from callus. Figure 3C show root formation, Figure 3D shows a complete plantlet and Figure 3E shows a one month old plantlet in pot soil.

Figure 4 is a map of pBI121 with an 800 bp HindIII-BamHI fragment containing the cauliflower mosaic virus (CaMV) 35S promoter cloned upstream of the GUS gene.

Figures 5A-C demonstrate rejuvenation of a tree. Figure 5A shows a two-year old tree. Figure 5B shows adventitious bud induction. Figure 5C shows propagated adventitious buds with plumules.

Figure 6A shows adventitious buds and Figure 6B shows stem pieces as explants for transformation.

Figure 7 demonstrates selection and induction of putative transgenic adventitious buds.

Figures 8A and 8B illustrate GUS staining of adventitious buds after selection for 5 months.

Figures 9A-C illustrate GUS staining of young transgenic stem pieces. Figures 9A-B show stem pieces and Figure 9C shows a shoot.

Figures 10A-C show GUS staining of transgenic leaf and leaf pieces.

Figure 11 shows the results of a Southern blot to a NPTII probe. 20 µg of DNA was digested with HindIII, run on a gel and blotted. Lanes 1-6 are transgenic lines. Lane ck is a negative control using DNA from a non-transgenic plant. Lane 7 is a positive control using DNA from a tomato transgenic line by plasmid pWS42 with NPTII as the selection marker.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to methods for genetically transforming tissue of trees, especially *Acacia mangium*, and regenerating whole plants via organogenesis from the transformed tissue.

The present invention is further detailed in the following Examples, which are offered by way of illustration and are not intended to limit the invention in any manner. Standard techniques well known in the art or the techniques specifically described below are utilized.

EXAMPLE 1

Reagents and Culture Media

A. MS basic medium: MS medium (Murashige and Skoog, 1962)

Name	Molecular formula	Concentration (mg/L)	Sigma Cat. No.
<u>Macronutrients</u>			
Ammonium Nitrate	NH ₄ NO ₃	1,650	A-3795
Potassium Nitrate	KNO ₃	1,900	P-8291
Calcium Chloride dihydrate	CaCl ₂ ·2H ₂ O	440	C-2536
Magnesium Sulfate Heptahydrate	MgSO ₄ ·7H ₂ O	370	M-7774
Potassium phosphate monobasic, anhydrous	KH ₂ PO ₄	170	P-8416
Ferrous Sulfate Heptahydrate	FeSO ₄ ·7H ₂ O	27.8	F-8263
Ethylenediamine-tetraacetic acid (EDTA)	C ₁₀ H ₁₄ N ₂ O ₈ Na ₂ · 2H ₂ O (Na ₂ EDTA)	37.3	E-6635

<u>Micronutrients</u>			
Potassium Iodide	KI	0.83	P-8166
Boric Acid	H ₃ BO ₃	6.2	B-9645
Manganese Sulfate monohydrate	MnSO ₄ ·H ₂ O	16.9	M-7899
Zinc Sulfate	ZnSO ₄ ·7H ₂ O	8.6	Z-1001
Molybdic Acid (Sodium salt: dihydrate)	Na ₂ MoO ₄ ·2H ₂ O	0.25	M-1651
Cupric Sulfate (Pentahydrate)	CuSO ₄ ·5H ₂ O	0.025	C-3036
Cobalt Chloride (hexahydrate)	CoCl ₂ ·6H ₂ O	0.025	C-2911
<u>Organic Reagents</u>			
myo-Inositol	C ₆ H ₁₂ O ₆	100	I-3011
Nicotinic acid	C ₆ H ₅ NO ₂	0.5	N-0765
Glycine	C ₂ H ₃ NO ₂	2.0	G-6143
Thiamine (Vitamin B1)	C ₁₂ H ₁₇ ClN ₄ OS·HCl	0.1	T-3902
Pyridoxine (Vitamin B6) Hydrochloride	C ₈ H ₁₁ NO ₃ ·HCl	0.5	P-9755

B. Other Organic Reagents

Name	Molecular formula	Sigma Cat.
		No.
L-Ascorbic acid (Vitamin C)	C ₆ H ₈ O ₆	A-2174
Casein Enzymatic Hydrolysate (CH)		C-7290
L-Glutamine (Gln)	C ₅ H ₁₀ N ₂ O ₃	G-9273
L-Asparagine monohydrate (Asn)	C ₄ H ₈ N ₂ O ₃ ·H ₂ O	A-4284
L-Proline (Pro)	C ₅ H ₉ NO ₂	P-4655
Sucrose		S-5390

C. Plant Growth Regulators

Name	Sigma Cat.
	No.
Indole-3-acetic Acid (IAA)	I-2886
α -Naphthaleneacetic Acid(NAA)	N-0640
1-Phenyl-3-(1,2,3-thiadiazol-5-yl)Urea (Thidiazuron, TDZ)	P-6186
6-Benzylaminopurine (6-BA)	B-3408
Kinetin (KT)	K-0753
Gibberellic Acid (GA ₃)	G-7645

D. Agar

Name	Sigma Cat.
	No.
Type M	A-4800
Purified	A-7921
Phytigel	P-8169

E. Antibiotics.

Timentin (T)	Beecham Pharmaceuticals (Pte) Ltd
Kanamycin (K)	Sigma
Genetins (G418 Sulfate)	Clontech 8056-2

F. Media for callus induction

AM-5	MS + 2,4-D 2.0 mg/L + KT 3.0 mg/L
AM-6	MS + 2,4-D 2.0 mg/L + KT 0.5 mg/L
AM-7	MS + NAA 2.0 mg/L + KT 3.0 mg/L
AM-8	MS + NAA 2.0 mg/L + KT 0.5 mg/L
AM-14	MS + 2,4-D 0.5 mg/L + 6-BA 3.0 mg/L
AM-15	MS + 2,4-D 1.0 mg/L + 6-BA 3.0 mg/L
AM-16	MS + 2,4-D 2.0 mg/L + 6-BA 3.0 mg/L
AM-17	MS + 2,4-D 0.5 mg/L + 6-BA 0.5 mg/L
AM-18	MS + 2,4-D 1.0 mg/L + 6-BA 0.5 mg/L
AM-19	MS + 2,4-D 2.0 mg/L + 6-BA 0.5 mg/L
AM-20	MS + 2,4-D 0.5 mg/L + 6-BA 1.0 mg/L
AM-21	MS + 2,4-D 1.0 mg/L + 6-BA 1.0 mg/L
AM-22	MS + 2,4-D 2.0 mg/L + 6-BA 1.0 mg/L
AM-27	MS + NAA 0.5 mg/L + 6-BA 3.0 mg/L
AM-28	MS + NAA 1.0 mg/L + 6-BA 3.0 mg/L
AM-29	MS + NAA 2.0 mg/L + 6-BA 3.0 mg/L
AM-30	MS + NAA 0.5 mg/L + 6-BA 0.5 mg/L
AM-31	MS + NAA 1.0 mg/L + 6-BA 0.5 mg/L
AM-32	MS + NAA 2.0 mg/L + 6-BA 0.5 mg/L
AM-33	MS + NAA 0.5 mg/L + 6-BA 1.0 mg/L
AM-34	MS + NAA 1.0 mg/L + 6-BA 1.0 mg/L
AM-35	MS + NAA 2.0 mg/L + 6-BA 1.0 mg/L
AM-231	MS + 2,4-D 5.0 mg/L + KT 0.5 mg/L
AM-233	MS + 2,4-D 0.5 mg/L + KT 1.0 mg/L
AM-234	MS + 2,4-D 1.0 mg/L + KT 1.0 mg/L

G. Media for adventitious buds induction

- a. AM-265: MS basic medium with TDZ 1.0 mg/L, IAA 0.25 mg/L, CH 100 mg/L, Vc 100 mg/L, Gln 150 mg/L, Asn 150 mg/L and Pro 150 mg/L, pH 5.8 after autoclaving, phytigel 0.275 or 0.30%, or agar 0.8% (A-4800, Sigma), sucrose 30 g/L.
- b. AM-261: MS basic medium with TDZ 1.0 mg/L, IAA 0.5 mg/L, CH 100 mg/L, Vc 100 mg/L, Gln 150 mg/L, Asn 150 mg/L, Pro 150 mg/L, pH 5.8 after autoclave, phytigel 0.275 or 0.30%, or agar 0.8% (A-4800, Sigma), sucrose 30 g/L.
- c. AM-304: MS basic medium with TDZ 2.0 mg/L, IAA 0.25 mg/L, CH 100 mg/L, Vc 100 mg/L, Gln 150 mg/L, Asn 150 mg/L and Pro 150 mg/L, pH 5.8 after autoclave, phytigel 0.275 or 0.30%, or agar 0.8% (A-4800, Sigma), sucrose 30 g/L.
- d. AM-308: MS basic medium with TDZ 1.0 mg/L, IAA 2.0 mg/L, CH 100 mg/L, Vc 100 mg/L, Gln 150 mg/L, Asn 150 mg/L and Pro 150 mg/L, pH 5.8 after autoclaving, phytigel 0.275 or 0.30%, or agar 0.8% (A-4800, Sigma), sucrose 30 g/L.

H. Media for pinnate leaf formation

- a. AM-337: MS basic medium with TDZ 0.01 mg/L, CH 100 mg/L, Vc 100 mg/L, Gln 150 mg/L, Asn 150 mg/L and Pro 150 mg/L, pH 5.8 after autoclaving, phytigel 0.275 or 0.30%, or agar 0.8% (A-4800, Sigma), sucrose 30 g/L.
- b. AM-41: MS basic medium supplemented with 6-BA 2 mg/L, CH 100 mg/L, Vc 100 mg/L, Gln 150 mg/L, Asn 150 mg/L and Pro 150 mg/L, pH 5.8 after autoclaving, phytigel 0.275 or 0.30%, or agar 0.8%, sucrose 30 g/L.

I. Media for root formation

- a. Modified AM-8: MS basic medium with NAA 2.0 mg/L, KT 0.5 mg/L, CH 100 mg/L, Vc 100 mg/L, Gln 150 mg/L, Asn 150 mg/L and Pro 150 mg/L, pH 5.8 after autoclaving, phytigel 0.30%, sucrose 30 g/L.
- b. AM-357: 1/2 MS basic medium with NAA 2.0 mg/L, KT 0.5 mg/L, CH 100 mg/L, Vc 100 mg/L, Gln 150 mg/L, Asn 150 mg/L and Pro 150 mg/L, pH 5.8 after autoclaving, phytigel 0.30%, sucrose 30 g/L.
- c. AM-451: 1/2 MS basic medium with NAA 2.0 mg/L, KT 0.1 mg/L, CH 100 mg/L, Vc 100 mg/L, Gln 150 mg/L, Asn 150 mg/L and Pro 150 mg/L, pH 5.8 after autoclaving, phytigel 0.35%, sucrose 20 g/L.

J. Transplantation

Peat soil:white sand 3:1

K. Media for *Agrobacterium tumefaciens* activationLB medium

Tryptone	1.0%
Yeast extract	0.5%
NaCl	0.8%

pH 7.0 before autoclaving (Sambrook et al., 1989)

YEP medium (Per liter) (Chilton et al., 1974)

Bactopeptone	10 g
Yeast extract	10 g
NaCl	5 g

AB medium

20 x Phosphate Buffer (per liter)

K_2HPO_4	60 g
NaH_2PO_4	20 g

Autoclave this solution separately

20 x Salts solution (per liter)

NH_4Cl	20 g
$MgSO_4 \cdot 7H_2O$	6 g
KCl	3 g
$CaCl_2$	0.2 g
$FeSO_4 \cdot 7H_2O$	0.05 g

pH to 7.0 before autoclaving

To make up the final medium, combine (to a final volume of 1 liter):

50 mL 20 x Phosphate Buffer

50 mL 20 x Salts Solution

900 mL sterile ddH₂O

Induction medium

MES buffer, pH 6.0 30 mM

1×AB medium

Glucose 0.5%

Acetosyringone 100 μM (a stock solution of acetosyringone must be made up fresh in DMSO)

Example 2Assay for GUS Activity

Stain the tissue overnight at 37°C in GUS staining solution. Gus staining solution is described by Jefferson (1987). It is X-gluc - 1 mM, sodium phosphate (pH 7.0) - 100 mM, EDTA - 10 mM, and Triton X-100 - 0.1%. GUS staining showed a positive blue reaction in adventitious buds and stem and leaf (Figures 8, 9, and 10).

Example 3Southern Analysis

A method for performing Southern blots is described in Sambrook et al. (1989). The method is:

A. DNA extraction

2-5 g fresh sample was frozen in liquid N₂. This was ground with a mortar and pestle in liquid nitrogen to a fine powder. The powder was transferred to a centrifuge tube (50 mL). 15 mL of Extraction Buffer was added, 2 mL 10% SDS was added and mixed thoroughly. This was incubated at 65°C for 15 minutes. 5 mL 5 M KAc was added and shaken vigorously. The mixture was incubated in ice for 20 minutes and then spun at 25,000 x g for 20 minutes. The supernatant was filtered through Microcloth into a new tube. The DNA was precipitated with ½ volume of isopropanol, mixed and incubated at -20°C for 30 minutes.

The DNA was pelleted at 25,000 x g for 30 minutes, the supernatant was poured off, and the tube was inverted and air dried for 30 minutes. The pellet was dissolved with 0.7 mL of 1×TE (pH 8.0) and transferred to an Eppendorf tube. This was spun 10 minutes. The supernatant was transferred to a new tube, 7 µL RNase (10 mg/mL) was added and left at room temperature for 10 minutes, then 75 µL 3 M NaAc (pH 5.3) and 500 µL of isopropanol were added. The solution was mixed and the DNA was pelleted in a microcentrifuge at full speed for 5 minutes. The pellet was washed with 70% ethanol, air dried and dissolved with 100 µL 1×TE (pH 8.0).

B. Extraction buffer

100 mM Tris-HCl, pH 8	50 mL
50 mM EDTA, pH 8	50 mL
500 mM NaCl	50 mL
10 mM β -ME	0.6 mL
ddH ₂ O	to 500 mL

C. Enzyme digestion

Reaction system I: DNA sample 100 μ L (20 μ g), 10 \times HindIII buffer 40 μ L, HindIII 8 μ L (80 units), add sterile double distilled H₂O 252 μ L to total volume 400 μ L.

The reaction was incubated at 37°C for overnight. 40 μ L of 3 M NaAc (pH 5.3) and 2/3 volume of 100% ethanol were added to the reaction system and this was incubated at -20°C for 30 minutes. This was spun at full speed at 4°C for 20 minutes. The supernatant was poured off and the tube was air dried for 30 minutes then the pellet was dissolved in 30 μ L sterile double distilled water.

D. Electrophoresis

Electrophoresis was performed on a 0.8% agarose gel in 1 \times TBE at 28V for overnight.

E. Transfer of DNA from Agarose gels to Nylon Membranes

This step was performed as described in Sambrook et al. (1989).

F. Prehybridization

A prehybridization solution of 6 \times SSC, 5 \times Denhardt's reagent, 0.5% SDS, 100 μ g/mL denatured, fragmented salmon sperm DNA (Stratagene products) and 50% formamide was prepared. 50 \times Denhardt's is: 5 g of Ficoll, 5 g of polyvinylpyrrolidone, 5 g of bovine serum albumin and ddH₂O to 500 mL, filtered and stored at -20°C.

After fixing DNA to the membrane, the membrane was placed into a hybridization tube containing suitable prehybridization solution using 0.2 mL prehybridization solution for each square centimeter of nylon membrane. The membrane was incubated at 42°C for 6 hours.

G. Probe label

During the prehybridization, labeled probe was prepared using a Boehringer Mannheim-High Primer DNA Labeling Kit. 50 ng of NPTII were added then brought to a volume of 8 μ L with H₂O. The DNA was denatured in a 100°C heat block for 10 minutes, chilled quickly in ice, and pulse spun.

On ice the denatured DNA was mixed with: 4 μ L High Prime reaction mixture, 3 μ L of dATP, dGTP, dTTP mixture, and 5 μ L of α -³²P dCTP, 3000 Ci/mmol (Biolab). This was incubated at 37°C for 10 minutes. The reaction was stopped by adding 20 μ L of 50 mM EDTA (pH 8.0).

The labeled probe was purified by running through a small Sephadex G50 column prepared on a small Pasteur pipette. The eluent was monitored with a counter and the first peak was collected.

H. Hybridization

The probe was added into the hybridization tube, then incubated at 42°C for 10-24 hours.

I. Washing membrane

The hybridization solution was poured off and the membrane was submerged in 2×SSC, 0.5% SDS at room temperature for 10 minutes. The membrane was transferred into 2×SSC, 0.1% SDS at room temperature for 15 minutes. The solution was replaced with 0.5×SSC, 0.1% SDS and the membrane was incubated at room temperature for 15 minutes. The solution was replaced with 0.1×SSC, 0.1% SDS and incubation continued at 55°C for 30-55 minutes. The membrane was transferred into 0.1×SSC at room temperature for 3-5 minutes, then air dried on 3MM Whatman paper for 30 minutes.

J. Autoradiography

The membrane was exposed to X-ray film (Kodak) to obtain an autoradiographic image at -80°C for one day or more.

Using NPTII fragment as a probe, Southern blotting showed that the NPTII gene had integrated in adventitious buds (Figure 11). The results demonstrate that this protocol of *Acacia mangium* transformation is very successful.

EXAMPLE 4Regeneration of *A. mangium*

A. Embryo culture

Mature seeds (black coat) were pre-treated with 98% H_2SO_4 for 2-3 minutes and washed with tap water several times. Treated seeds were sterilized with 70% ethanol for 2-3 minutes and washed 5 times with sterile ddH_2O . Seeds were then immersed in 0.1% HgCl_2 for 6 minutes and washed 5 times with sterile ddH_2O , again sterilized with bleach 30% (market product) for 6 minutes, then washed 5 times with sterile ddH_2O . Sterilized seeds were soaked in sterile ddH_2O overnight for isolating zygotes for embryo culture. MS basic medium (Murashige and Skoog, 1962), pH 5.8, sucrose 30 g/L, phytigel 0.25% or agar 0.7% (Sigma, A-4800), with or without activated charcoal, was used to culture mature embryo. Isolated mature zygotic embryos were cultured on MS using a photoperiod of 12/12 hours or 16/8 hours (L/D) or complete dark, at 25-28°C. Germinated hypocotyls or leaves or petioles or stems were used as explants to induce callus formation.

B. Callus induction

Different media were used to induce callus, including medium Nos. AM-5, 6, 7, 8, 14, 15, 16, 17, 18, 19, 20, 21, 22, 27, 28, 29, 30, 31, 32, 33, 34, 35, 231, 233 and 234 at 12/12 hours or 16/8 hours (L/D) or Dark at 25°C. Callus induced from young leaflet: AM-L; from young petiole: AM-P; from hypocotyl: AM-H; from young stem: AM-S; from bud: AM-B; from root: AM-R. All above calli were used to induce adventitious buds, using a photoperiod of 16/8 hours, 1800-2000 lux, and 28°C. The results of callus induction on different media were different (Table.1). Generally callus induction was not difficult. All explants produced calli at the photoperiod of 12/12 hrs or 16/8 hrs (L/D) (see, e.g., Figure 1A; Figure 2A; Figure 3A), and 25°C for 20 days.

Table 1

Results of callus induction on different media
at 12/12 hours (L/D), 25°C cultured for 35 days

Medium No.	Number of Explants (leaflet)	No. of explants forming callus	No. of explants complete callus formation
AM-5	40	40	33
AM-6	40	40	31
AM-7	40	40	3
AM-8	40	40	4
AM-14	40	40	36
AM-15	40	40	40
AM-16	40	40	38
AM-17	40	40	36
AM-18	40	40	39
AM-19	40	40	39
AM-20	40	40	38
AM-21	40	40	40
AM-22	40	40	40
AM-27	40	40	12
AM-28	40	40	8
AM-29	40	40	11
AM-30	40	40	10
AM-31	40	40	6
AM-32	40	40	2
AM-33	40	40	4
AM-34	40	40	4
AM-35	40	40	6
AM-231	40	40	30
AM-233	40	40	20
AM-234	40	40	20
	Petiole		
AM-6	20	20	15
AM-17	20	20	20

C. Adventitious bud induction

Calli induced from leaflet, petiole, young stem and buds were cultured on AM-261, AM-265, AM-304, and AM-308, at a photoperiod of 16/8 hrs (L/D), at 28°C. One month later, there were some bud recoveries from callus (Figure 1B; Figure 2B). Bud recovery from friable callus first appeared as compact and smooth bud terminal. The ratio of bud recovery reaches 15% on AM-265. But on these media, induced buds could not easily form pinnate leaves and elongate.

D. Pinnate leaf formation and adventitious bud elongation

Bud recoveries from callus were transferred into AM-337 or AM-41 with GA₃ 2.5 mg/L, at a photoperiod of 16/8 hrs (L/D), 1800-2000 lux, and 28°C. After culture for one month, some of them could form pinnate leaves and elongate (Figures 1C, D, E; Figure 2C; and Figure 3B). The ratio of pinnate leaf formation and bud elongation reaches 16.67%.

E. Root formation

Elongated buds were cultured in AM-357 or AM-451 for root formation. 20 days later, adventitious roots began to appear from basic stem of buds (Figure 3C). After root formation, plantlets were transferred to MS basic medium without any plant growth regulators. Plantlets grow normally and fast, and the root system grew well (Figure 3D).

F. Transplantation

Phytigel on plantlets was washed away and the plantlets were transferred to pot (peat soil:white sand 3:1) and grown in a growth chamber at a photoperiod of 16/8 hours and 25°C. One month later the plantlets were transferred to a green house (Figure 3E).

Example 5

Rejuvenation of Tree

The auxiliary buds of a two-year old young tree (Figure 5A) were cultured on MS basic medium with NAA 0.1 mg/L, 6-BA 3.0 mg/L, CH 100 mg/L, Vc 100 mg/L, Pro 150 mg/L, Asn 150 mg/L and Gln 150 mg/L, phytigel 0.275% or agar 0.8% (A-4800, Sigma), pH 5.8 after autoclaving at 121°C, sucrose 30 g/L. Sixty days later, some adventitious buds with phyllodes were obtained (Figure 5B). Induced adventitious buds with phyllodes were subcultured on AM-41 (MS basic medium with 6-BA 2 mg/L, CH 100 mg/L, Vc 100 mg/L, Pro 150 mg/L, Asn 150 mg/L and Gln 150 mg/L, phytigel 0.275% or agar 0.8% (A-4800, Sigma), pH 5.8 after autoclaving at 121°C, sucrose 30 g/L). After sub-culturing adventitious buds with phyllodes twice in about two months, the rejuvenation of adventitious shoots with pinnate leaf was obtained (Figure 5C). Adventitious buds can be used as explants for transformation (Figure 6A).

Example 6

A. mangium Genetic Transformation

A. Activation of *Agrobacterium tumefaciens* Strain LBA4404/ pBI121

Agrobacterium strain LBA4404 (Ooms et al., 1981) harbouring binary vector plasmid pBI121 (size of vector 13.0 kb, Clontech, Figure 4) was used for this experiment. pBI121 was derived from pBI101 with an 800-bp HindIII-BamHI fragment containing the cauliflower mosaic virus (CaMV) 35S promoter cloned upstream of the GUS gene. From a permanent glycerol stock stored at -70°C, *Agrobacterium tumefaciens* pBI121/LBA4404 was streaked onto a solid LB (pH 7.0) medium plate containing Streptomycin 100 mg/L (Str¹⁰⁰) and Kanamycin 50 mg/L (K⁵⁰). This was incubated 2-3 days at 28°C. Fresh pBI121/LBA4404 was streaked onto solid LB medium (pH 7.0) with Streptomycin 100 mg/L (Str¹⁰⁰) and Kanamycin 50 mg/L (K⁵⁰) for overnight or 24 hours at 28°C in the dark. Several colonies of pBI121/LBA4404 were picked and pBI121/LBA4404 was inoculated in liquid 50 mL LB medium with Streptomycin 100 mg/L (Str¹⁰⁰) and Kanamycin 50 mg/L (K⁵⁰), and cultured at 28°C under dark, 250 rpm for 10 hours until OD₆₀₀ = 0.70-1.10. This was spun down at 3500 rpm for 30 minutes or 5000 rpm for 10 minutes, and resuspended in 4-5 volumes of YEP medium (pH 7.0) with Streptomycin 100 mg/L (Str¹⁰⁰) and Kanamycin 50 mg/L (K⁵⁰), OD₆₀₀ = 0.10-0.20, incubated at 28°C in the dark, 250 rpm for 8-10 hours, OD₆₀₀ = 0.70-1.20. This was spun down at 3500 rpm for 30 minutes and resuspended in the same volume of sterile NaCl 0.9%. This was spun down at 3500 rpm for 30 minutes and resuspended in 2-3.5 volumes of induction medium, OD₆₀₀ = 0.2-0.3, and incubated at 28°C in the dark, 120 rpm for 8-15 hours, OD₆₀₀ = 0.70-1.20 for infecting explants or callus or a cell suspension.

B. Preculture of stem pieces

Stem pieces were cultured on AM-265 for 3 days using a photoperiod of 16/8 hrs (L/D), 1800-2000 lux, at 28°C.

C. Co-cultivation

After preculture, stem pieces were soaked in 0.5 M mannitol for 20-25 minutes and then transferred to activated pBI121/LBA4404 suspension for 15 minutes. Infected young stem pieces

were washed once and dried on sterile Whatman paper before being cultured on AM-265 with 100 μ M acetosyringone at pH 5.2 at 22°C in the dark for 3 days.

D. Selection of transformed adventitious buds and transformed callus

After co-culturing for 3 days, stem pieces were washed with sterile ddH₂O for 10 times and dried on sterile Whatman paper. Then stem pieces were cultured on AM-265 containing Timentin 250 mg/L and G418 12 mg/L, phytigel 0.275%, using a photoperiod of 16/8 hrs (L/D) and 28°C for selecting transformed callus or transformed adventitious buds. Subsequently, stem pieces were cultured on the above media with G418 12 mg/L for 25 days, G418 20 mg/L for 60 days, G418 30 mg/L for 25 days, then on G418 12 mg/L. After continuous selecting for 4 months on antibiotics, 33.75% of stem pieces formed adventitious buds, and 2.5 mg/L to 5 mg/L GA₃ was added into the same media to promote adventitious bud elongation (Figure 7). After selection for five months, Timentin was no longer used in the medium and some adventitious buds were used for GUS staining. GUS staining showed positive blue colour reaction in adventitious buds (Figure 8, Figure 9, and Figure 10). Adventitious buds were transferred to AM-41 with GA₃ 2.5 mg/L to promote pinnate leaf formation.

E. Root formation

Transformed adventitious buds were transferred into AM-357 or AM-451 with or without G418 (10 mg/L) using a photoperiod of 16/8 hrs (L/D) at 28°C.

While the invention has been disclosed in this patent application by reference to the details of preferred embodiments of the invention, it is to be understood that the disclosure is intended in an illustrative rather than in a limiting sense, as it is contemplated that modifications will readily occur to those skilled in the art, within the spirit of the invention and the scope of the appended claims.

List of References

- Ahmad DH (1991). "Micropropagation of *Acacia mangium* from aseptically germinated seedlings" *Journal of Tropical Forest Science* 3(3):204-208.
- Anwar C (1992). "The growth of Shorea seedlings on soil media of several age levels of *Acacia mangium* stands" *Buletin Penelitian Hutan* 0(544):9-16.
- Awang K (1994). "Growth of three multipurpose tree species on tin tailing in Malaysia" *Journal of Tropical Forest Science* 7(1):106-112.
- Basu PK, Ganguly DK, and Mandal GS (1987). "Introduction of exotics in south-west Bengal (India): *Acacia mangium* in coastal area (Digha-Midnapur)" *Indian Forester* 113(10):675-680.
- Bayliss M (1998a). *Asia Pacific Papermaker* 8(1):19.
- Bayliss M (1998b). *Asia Pacific Papermaker* 8(1):64.
- Bhaskan P and Subbash K (1996). "Micropropagation of *Acacia mangium* Willd. through nodal bud culture" *Indian Journal of Experimental Biology* 34:590-591.
- Chilton MD, Currier TC, Farrand SK, Bendich AJ, Gordon MP and Nester EW (1974). "*Agrobacterium tumefaciens* DNA and PS8 bacteriophage DNA not detected in crown gall tumors" *Proc. Natl. Acad. Sci. USA* 71:3672-3676.
- Cole TG, Yost RS, Kablan R and Olsen T (1996). "Growth potential of twelve *Acacia* species on acid soils in Hawaii" *Forest Ecology and Management* 80(1-3):175-186.
- Duguma B, Tonye J, Kanmegne J, Manga T and Enoch T (1994). "Growth of ten multipurpose tree species on acid soils in Sangmelima, Cameroon" *Agroforestry Systems* 27(2):107-119.
- Duke JA (1984). "*Acacia mangium* Willd", Handbook of Energy Crops (unpublished).
- Galiana A, Tibok A and Duhoux E (1991a). "*In Vitro* propagation of the nitrogen-fixing tree-legume *Acacia mangium* Willd" *Plant and Soil* 135:151-159.
- Galiana A, Tibok A and Duhoux E (1991b). "Nitrogen-fixing potential of micropropagated clones of *Acacia mangium* inoculated with different *Bradyrhizobium* spp. strains" *Plant and Soil* 135(2): 161-166.
- Gutteridge RC and Shelton HM (1994). "Forage Tree Legumes" in Tropical Agriculture by CAB International.
- Jefferson RA (1987). "Assaying chimeric in plants: The GUS gene fusion system" *Plant Mol. Biol. Rep.* 5:387-405.

- Khemnark C (1994). "Rehabilitation of degraded tropical forest through agroforestry practices: A case study in Thailand" *Journal of Tropical Forest Science* 7(1):128-135.
- Latif MA, Das S, Rahman MF and Habib MA (1995). "Growth and yield tables for *Acacia mangium* in the plantation in Bangladesh" *Journal of Tropical Forest Science* 7(4):591-598.
- Majid NM, Hashim A and Abdol I (1994). "Rehabilitation of ex-tin mining land by agroforestry practice" *Journal of Tropical Forest Science* 7(1):113-127.
- Murashige T and Skoog F (1962). "A revised medium for rapid growth and bioassays with tobacco tissue cultures" *Physiol. Plant* 15:473-497.
- Ooms G, Hooykaas PJ, Moolenaar G and Schilperoort GA (1981). "Grown gall plant tumors of abnormal morphology, induced by *Agrobacterium tumefaciens* carrying mutated octopine Ti plasmids; analysis of T-DNA functions" *Gene* 14:33-50.
- Robinson C (1999). "Making forest biotechnology a commercial reality" *Nature Biotechnology* 17: 27-30.
- Sambrook J, Fritsch EF and Maniatis T (1989). Molecular Cloning - A Laboratory Manual (second edition), Cold Spring Harbor Laboratory Press.
- Simmons MH (1987). Growing Acacias (edited by Simmons MH) published by Kangaroo Press Pty Ltd.
- Tzfira T, Zuker A and Altman A (1998). "Forest-tree biotechnology: genetic transformation and its application to future forests" *TIBTECH* 16:439-446.
- Wibowo A, Suharti M and Pickford SG (1992). "Fuel characteristics and fire behaviour in alang-alang under *Acacia mangium* plantation in Depok, West Java" *Buletin Penelitian Hutan* 0(544):1-7.
- Widiarti A and Alrasjid H (1987). "Introduction of fuelwood trees species on degraded lands in Paseh and Kadipaten areas (Indonesia)" *Buletin Penelitian Hutan* 0(488):1-17.
- Williams ER, Gunn B, Reynolds D and Westcott M (1992). "Germination tests for small collection of *Acacia* seed" *Seed Science and Technology* 20(2):321-326.
- Wong WC, Ho KS and Wong CN (1988). "*Acacia mangium* from Sabah for plywood and decorative panel manufacture: Initial trials" *Journal of Tropical Forest Science* 1(1):42-50.
- Yusoff MNM, Chew LT, Ali ARM and Nasir NM (1989). "The adhesive properties of bark extract of *Acacia mangium*" *Journal of Tropical Forest Science* 2(2):104-109.

WHAT IS CLAIMED IS:

1. A method for regenerating *Acacia mangium* comprising:
 - a) inducing callus formation from an explant;
 - b) culturing said callus to produce adventitious buds;
 - c) culturing said adventitious buds to elongate and produce pinnate leaves; and
 - d) culturing elongated buds of step (c) such that they produce roots and become plantlets.
2. The method of claim 1 wherein seeds are cultured to produce said explant.
3. The method of claim 1 wherein said explant is selected from the group consisting of hypocotyls, cotyledons, leaves, petioles and stems.
4. The method of claim 1 wherein said explant is cultured on a medium comprising MS medium supplemented with an auxin and a cytokinin.
5. The method of claim 4 wherein said auxin is present at 0.5-2.0 mg/L and said cytokinin is present at 0.5-3.0 mg/L.
6. The method of claim 4 wherein said auxin is selected from the group consisting of 2,4-D and α -naphthaleneacetic acid and wherein said cytokinin is selected from the group consisting of kinetin and 6-benzylaminopurine.
7. The method of claim 1 wherein said callus is cultured on a medium comprising MS basic medium supplemented with a) thidiazuron, b) indole acetic acid, c) casein enzymatic hydrolysate, d) L-ascorbic acid, e) L-glutamine, f) L-asparagine, g) L-proline, h) sucrose and i) agar or phytagel.
8. The method of claim 1 wherein said adventitious buds are cultured on a medium comprising MS medium supplemented with a) thidiazuron, b) casein enzymatic hydrolysate, c) L-ascorbic acid, d) L-glutamine, e) L-asparagine, f) L-proline, g) sucrose and h) agar or phytagel.

9. The method of claim 1 wherein said elongated buds are cultured on a medium comprising 1/2 MS basic medium supplemented with a) α -naphthaleneacetic acid, b) kinetin, c) casein enzymatic hydrolysate, d) L-ascorbic acid, e) L-glutamine, f) L-asparagine, g) L-proline, h) sucrose and i) phytigel.
10. The method of claim 1 wherein said explant has been transformed.
11. A method for regenerating *Acacia mangium* comprising:
 - a) culturing auxiliary buds from an *Acacia mangium* tree to produce adventitious buds comprising phyllodes;
 - b) subculturing said adventitious buds comprising phyllodes to produce adventitious shoots;
 - c) culturing said adventitious shoots.
12. The method of claim 11 wherein said culturing of auxiliary buds is performed on a medium comprising MS basic medium supplemented with a) α -naphthaleneacetic acid, b) 6-benzylaminopurine, c) casein enzymatic hydrolysate, d) L-ascorbic acid, e) L-proline, f) L-asparagine, g) L-glutamine, h) sucrose, and i) phytigel or agar.
13. The method of claim 11 wherein said subculturing of adventitious buds comprising phyllodes is performed on a medium comprising MS basic medium supplemented with a) 6-benzylaminopurine, b) casein enzymatic hydrolysate, c) L-ascorbic acid, d) L-glutamine, e) L-asparagine, f) L-proline, g) sucrose and h) phytigel or agar.
14. A method of transforming *Acacia mangium* with a gene of interest comprising the steps of:
 - a) activating *Agrobacterium tumefaciens* comprising said gene of interest to form activated *Agrobacterium tumefaciens*;
 - b) preculturing an explant of *Acacia mangium* to yield a precultured explant;
 - c) co-cultivating said activated *Agrobacterium tumefaciens* and said precultured explant to produce infected explants;

- d) culturing said infected explants to induce callus and adventitious buds; and
e) culturing said callus or adventitious buds on a selective medium.
15. The method of claim 14 wherein said preculturing of said explant is performed on a medium comprising MS basic medium supplemented with a) thidiazuron, b) indole-3-acetic acid, c) casein enzymatic hydrolysate, d) L-ascorbic acid, e) L-glutamine, f) L-asparagine, g) L-proline, h) sucrose and i) phytagel or agar.
16. The method of claim 14 wherein said explants are soaked in 0.5 M mannitol prior to the step of co-cultivating.
17. The method of claim 14 wherein said co-cultivating is performed on a medium comprising MS basic medium supplemented with a) thidiazuron, b) indole-3-acetic acid, c) casein enzymatic hydrolysate, d) L-ascorbic acid, e) L-glutamine, f) L-asparagine, g) L-proline, h) sucrose and i) phytagel or agar.
18. The method of claim 14 wherein said co-cultivating is performed in the dark.
19. The method of claim 14 wherein said selective medium comprises a medium comprising MS basic medium supplemented with a) thidiazuron, b) indole-3-acetic acid, c) casein enzymatic hydrolysate, d) L-ascorbic acid, e) L-glutamine, f) L-asparagine, g) L-proline, h) sucrose and i) phytagel or agar.
20. The method of claim 14 wherein said preculture is performed using a photoperiod of 16 hours light/8 hours dark.
21. The method of claim 14 wherein said culturing on selective medium is performed using a photoperiod of 16 hours light/8 hours dark.
22. The method of claim 14 wherein said explant is selected from the group consisting of stem, leaflet, petiole and bud.

23. A method for promoting elongation of transformed adventitious buds of *Acacia mangium* comprising transforming an *Acacia mangium* explant by the method of claim 14 and further comprising a step of addition of gibberellic acid to the culture medium following formation of adventitious buds.
24. A method for promoting pinnate leaf formation on transformed adventitious buds of *Acacia mangium* comprising transforming an *Acacia mangium* explant by the method of claim 14 and further comprising culturing adventitious buds which develop on a medium with gibberellic acid.
25. A method for promoting root formation from transformed adventitious buds comprising culturing transformed adventitious buds on a medium comprising 1/2 MS basic medium supplemented with a) α -naphthaleneacetic acid, b) kinetin, c) casein enzymatic hydrolysate, d) L-ascorbic acid, e) L-glutamine, f) L-asparagine, g) L-proline, h) sucrose and i) phytagel.
26. The method of claim 25 wherein said culturing is performed using a 16 hour light/8 hour dark photoperiod.
27. The method of claim 25 wherein said culturing is performed at 28°C.
28. A method of preparing transgenic *Acacia mangium* cells comprising the steps of
a) preculturing stem pieces of *Acacia mangium* in a culture medium; and
b) cocultivating said stem pieces of step (a) with activated *Agrobacterium tumefaciens*.
29. The method of claim 28 wherein said preculturing is performed for 3 days using a photoperiod of 16/8 hours (light/dark).
30. The method of claim 29 wherein said preculturing is performed using 1800-2000 lux for the light cycles.

31. The method of claim 28 wherein said preculturing is performed at 28°C.
32. The method of claim 28 wherein said culture medium is AM-265.
33. The method of claim 28 wherein said stem pieces are soaked in a mannitol solution prior to cocultivating with *Agrobacterium tumefaciens*.
34. The method of claim 28 wherein said activated *Agrobacterium tumefaciens* were prepared by growing them in induction medium.
35. The method of claim 34 wherein said activated *Agrobacterium tumefaciens* were prepared by growing them in the dark.
36. The method of claim 34 wherein said activated *Agrobacterium tumefaciens* were prepared by growing them at 28°C.
37. A method of making transgenic *Acacia mangium* plants comprising;
 - a) preparing transgenic *Acacia mangium* cells by the method of claim 28;
 - b) culturing said cells in a selective medium,
 - c) adding a growth promoter; and
 - d) rooting buds which develop.
38. The method of claim 37 wherein said selective medium comprises an antibiotic.
39. The method of claim 37 wherein said culturing is performed for more than 1 month.
40. The method of claim 37 wherein said growth promoter is gibberellic acid.
41. A transgenic *Acacia mangium* cell.
42. A transgenic *Acacia mangium* plant.

Fig. 1 Adventitious bud induction from leaf
A: Callus induction; B: Adventitious bud recovery;
C,D,E: Plumule formation and adventitious bud elongation





2/11




Fig.2 *Acacia mangium* regeneration from petiole. A: Callus induction; B: Adventitious bud recovery; C: Plumule formation and adventitious bud elongation.



3/11




Fig.3 *Acacia mangium* regeneration from stem pieces A: Callus; B: Adventitious buds induction from callus; C: Root formation ; D: Complete plantlet; E: One month old plantlet in pot soil.



pBI121 vector




Fig.4 Map of pBI121 with an 800-bp HindIII-BamHI fragment containing the cauliflower mosaic virus (CaMV) 35S promoter cloned upstream of the GUS gene. Vector size: 13.0kb(CLONTECH)



5/11





Fig.5 Rejuvenation of tree. A: Two-year old tree; B: Adventitious bud induction;
C: Propagated adventitious buds with plumules



6/11



B



A

Fig.6 A: Adventitious buds; B: Stem pieces as explants for transformation



7/11




Fig. 7 Selection and induction of putative transgenic adventitious buds



8/11



Fig. 8 Gus staining of adventitious buds after selection for 5 months



9/11

Fig.9 GUS staining of young transgenic stem pieces

A,B: stem pieces C: A shoot

A




B



C





11/11





Fig. 11 Southern blot to nptII probe: 20 μ g DNA was digested with Hind III and hybridized to nptII probe. 1-6, transgenic lines; ck, negative control: DNA from non-transgenic plant; 7, positive control: DNA from a tomato transgenic line by plasmid pWS42 with nptII as selection marker



10/11



Fig.10 GUS staining of transgenic leaf and leaf pieces

